

英文論文

- 1, Whiteside L.A and Nagamine R :
Biomechanical aspects of knee replacement designs
In : Scott W.N (ed) : The Knee - Chapter 59, .pp. 1079 - 1096,
Mosby-Year Book, Inc. St. Louis, 1993
- 2, Nagamine R, White S.E, McCarthy D.S and Whiteside L.A :
Patellar tracking after total knee arthroplasty. : The effect of tibial tray
malrotation and articular surface configuration.
Clin Orthop 1994;304: 263-271.
- 3, Nagamine R, Whiteside L.A, Otani T, White S.E and Milliano M.T :
Patellar tracking measurement in the normal knee.
J Orthop Res 1995;13: 115-122.
- 4, Nagamine R, White S.E and Whiteside L.A :
Effect of rotational malposition of the femoral component on knee stability
kinematics after total knee arthroplasty. J Arthroplasty 1995;10 : 265-270.
- 5, Nagamine R, Whiteside L.A., Otani T, White S.E and McCarthy D.S. :
Effect of medial displacement of the tibial tubercle on patellar position after
rotational malposition of the femoral component in total knee arthroplasty.
J Arthroplasty 1996;11: 104-110.
- 6, Miura H, Kawamura H, Nagamine R, Urabe K and Iwamoto Y.
Is patellar height really lower after high tibial osteotomy?
Fukuoka Acta Med. 1997;88(6) : 261-6.
- 7, Nagamine R, Miura H, Inoue Y, Tanaka K, Urabe K, Nishizawa M, Okamoto Y
and Iwamoto Y
Malposition of the tibial tubercle during flexion in knees with patellofemoral
arthritis. Skeletal Radiology 1997;26:597-601.
- 8, Nagamine R, Miura H, Inoue Y, Tanaka K, Urabe K, Nishizawa M, Okamoto Y
and Iwamoto Y
Reliability of the anteroposterior axis and the posterior condylar axis for
rotational alignment of the femoral component in total knee arthroplasty.
J orthop Sci. 1998;3(4):194-198.
- 9, Nagamine R, Whiteside LA, Miura H, Urabe K, Matsuda S and Iwamoto Y
Rotational malalignment of the femoral and tibial components in total knee
arthroplasty.
Joint Arthroplasty. Imura S, Wada M and Omori H. edited 373-383, 1999
Springer Tokyo

- 10, Nagamine R, Miura H, Urabe K, Matsuda S, Chen W. J, Matsunobu T and Iwamoto Y
Radiological assessment of the position of the tibial tuberosity by means of a marking wire in knees with patellofemoral arthritis.
Skeletal Radiology 1999;28:27-32.
- 11, Miura H, Kawamura H, Arima J, Mawatari T, Nagamine R, Urabe K and Iwamoto Y
A new, all-inside technique for meniscus repair.
Arthroscopy 1999;15(4):453-455.
- 12, Matsuda S, Miura H, Nagamine R, Urabe K, Ikenoue T, Okazaki K and Iwamoto Y
Posterior tibial slope in the normal and varus Knee.
Am J knee Surg. 1999;12(3):165-168.
- 13, Matsuda S, Miura H, Nagamine R, Urabe K, Harimaya K, Matsunobu T. and Iwamoto Y
Changes in knee alignment after total knee arthroplasty.
J Arthroplasty 1999;14: 566-570.
- 14, Matsuda S, Miura H, Nagamine R, Urabe K, Matsunobu T. and Iwamoto Y
Knee stability in posterior cruciate ligament retaining total knee arthroplasty.
Clin Orthop 1999;366: 169-173.
- 15, Nagamine R, Miura H, Bravo CV, Urabe K, Matsuda S, Miyanishi K, Hirata G and Iwamoto Y
Anatomic variations should be considered in total knee arthroplasty.
J Orthop Sci 2000;5(4):232-237.
- 16, Miura H, Noguchi Y, Mitsuyasu H, Nagamine R, Urabe K, Matsuda S, and Iwamoto Y
Clinical features of multiple epiphyseal dysplasia expressed in the knee.
Clin Orthop 2000;380:184-190.
- 17, Miyanishi K, Nagamine R, Murayama S, Miura H, Urabe K, Matsuda S, Hirata G and Iwamoto Y
Tibial tubercle malposition in patellar joint instability. -A computed tomography study in full extension and at 30° flexion-
Acta Orthop Scand 2000;71(3):286-291.
- 18, Nagamine R, Hanada Y, Kondo K, Fukumoto S, Shuto T, Nakashima Y, Hirata G, Katayama A and Iwamoto Y
Quantification of bone volume on radiographs using NIH Image.
Mod Rheumatol 2000;10:220-224.

- 19, Takeda T, Morimoto N, Kinukawa N, Nagamine R, Shuto T, Iwamoto Y and Miyahara H
Factors affecting emotional instability in female rheumatoid arthritis outpatients with limited functional disorder.
Mod Rheumatol 2000;10:240-246.
- 20, Takeda T, Morimoto N, Kinukawa N, Nagamine R, Shutou T and Iwamoto Y
Factors affecting depression and anxiety in female Japanese patients with rheumatoid arthritis.
Clin Exp Rheumatol 2000;18:735-738.
- 21, Matsuda S, Miura H, Nagamine R, Urabe K, Hirata G and Iwamoto Y
Effect of femoral and tibial component position on patellar tracking following total knee arthroplasty. 10-Year Follow-Up of Miller-Galante I Knees
Am J knee Surg. 2001;14(3):152-156.
- 22, Nagamine R, Miura H, Urabe K, Matsuda S, Hirata G, Moro-oka T, Kawano T and Iwamoto Y
A new concept for precise patella resection in total knee arthroplasty
Am J knee Surg. 2001;14(4):227-231.
- 23, Inoue S, Nagamine R, Miura H, Urabe K, Matsuda S, Sakaki K and Iwamoto Y
Anteroposterior weight-bearing radiography of the knee with both knees in semiflexion using new equipment. J Orthop Sci. 2001;6:475-480.
- 24, Nagamine R, Maeda T, Shuto T, Nakashima Y, Hirata G and Iwamoto Y
Menopausal syndrome in female patients with rheumatoid arthritis
Mod Rheumatol. 2001;11:230-233.
- 25, Jo Y, Matsumoto T and Nagamine R
A case of Crohn's disease with leg pain
Br J Radiol. 2001;74:203-204.
- 26, Maeda K, Yamada H, Nagamine R, Shuto T, Nakashima Y, Hirata G and Iwamoto Y
Involvement of CD4+, CD57+ T cells in the disease activity of rheumatoid arthritis
Arthritis Rheum. 2002;46(2):379-384.
- 27, Tsuchiya K, Nagamine R and Iwamoto Y
Discovertebral lesion in ankylosing spondylitis - Differential diagnosis with discitis by magnetic resonance imaging-
Mod Rheumatol. 2002;12:113-117.

- 28, Sakamoto A, Nagamine R, Maeda T, Shuto T, Hirata G, Nakashima Y, Matsuda S and Iwamoto Y
Multicentric reticulo-histiocytosis with hydrarthrosis in both knee joints: disease stabilization with synovectomy, and medication with a steroid and low-dose methotrexate
Mod Rheumatol. 2002;12:155-159.
- 29, Moro-oka T, Matsuda S, Miura H, Nagamine R, Urabe K, Kawano T, Higaki H and Iwamoto Y
Patellar tracking and patellofemoral geometry in deep knee flexion
Clin Orthop. 2002;394:161-168.
- 30, Kuwano T, Nagamine R, Urabe K and Iwamoto Y
New radiographic analysis of sesamoid rotation in Hallux Valgus: Comparison with conventional evaluation methods. Foot and Ankle, 2002;23(9):811-817.
- 31, Miyagi T, Matsuda S, Miura H, Nagamine R, Urabe K, Inoue S, Iwamoto Y
Changes in patellar tracking after total knee arthroplasty: 10-year follow-up of Miller-Galante I knees.
Orthopaedics, 2002;25(8):811-813.
- 32, Kawano T, Miura H, Nagamine R, Urabe K, Matsuda S, Mawatari T, Moro-oka T and Iwamoto Y
Factors affecting patellar tracking after total knee arthroplasty.
J Arthroplasty, 2002;17(7):942-947.
- 33, Matsuda S, Mizuuchi H, Miura H, Nagamine R, Urabe K and Iwamoto Y
Tibial shaft axis does not always serve as a correct coronal landmark in total knee arthroplasty for varus knees. J Arthroplasty, 2003;18:56-62.
- 34, Urabe K, Miura H, Kuwano T, Matsuda S, Nagamine R, Sakai S, Masuda K and Iwamoto Y
Comparison between the shape of resected femoral sections and femoral prostheses used in total knee arthroplasty in Japanese patients: simulation using three-dimensional computed tomography.
J Knee Surg. 2003;16(1):27-33.
- 35, Nagamine R, Miyanishi K, Miura H, Urabe K, Matsuda S and Iwamoto Y
Medial torsion of the tibia in Japanese patients with osteoarthritis of the knee
Clin Orthop, 2003;408:218-24.
- 36, Nabeyama R, Matsuda S, Miura H, Kawano T, Nagamine R, Mawatari T, Tanaka K, Iwamoto Y.
Changes in anteroposterior stability following total knee arthroplasty.
J Orthop Sci. 2003;8(4):526-31.

- 37, Kawano T, Miura H, Nagamine R, Urabe K, Matsuda S, Mawatari T, Moro-Oka T, Iwamoto Y.
Alignment in total knee arthroplasty following failed high tibial osteotomy.
J Knee Surg. 2003;16(3):168-72.
- 38, Matsuda S, Miura H, Nagamine R, Urabe K, Mawatari T, Iwamoto Y.
A comparison of rotational landmarks in the distal femur and the tibial shaft.
Clin Orthop. 2003;(414):183-8.
- 39, Nagamine R, Kondo K, Ikemura S, Shiranita A, Nakashima S, Ihara H,
and Sugioka S.
Distance of the femorotibial joint gap can be controlled in the modified
ligament dependent cut procedure during total knee arthroplasty.
Fukuoka Acta Med. 2003;94(10):296-303
- 40, Matsuda S, Miura H, Nagamine R, Mawatari T, Tokunaga M, Nabeyama R,
and Iwamoto Y.
Anatomical analysis of the femoral condyle in normal and osteoarthritic knees.
J Orthop Res. 2004;22(1):104-9.
- 41, Nagamine R, Kondo K, Ikemura S, Shiranita A, Nakashima S, Hara T, Ihara H,
and Sugioka Y.
Distal femoral cut perpendicular to the mechanical axis may induce varus
instability in flexion in medial osteoarthritic knees with varus deformity in total
knee arthroplasty. - A pitfall of the navigation system -
J Orthop Sci. 2004;9:555-9
- 42, Kuwano T, Urabe K, Miura H, Nagamine R, Matsuda S, Satomura M, Sasaki T,
Sakai S, Honda H, and Iwamoto Y
Importance of the lateral anatomic tibial slope as a guide to the tibial cut in
total knee arthroplasty in Japanese patients.
J Orthop Sci. 2005;10(1):42-47.
- 43, Nagamine R, Matsunobu T, Takayama M, Miura H, Matsuda S, and Iwamoto Y.
The extramedullary guide of the proximal tibia should be seen in straight in
front during total knee arthroplasty.
Fukuoka Acta Med. 2006;97(5):146-152.
- 44, Nagamine R, Inoue S, Miura H, Matsuda S, and Iwamoto Y.
Femoral shaft bowing influences the correction angle for high tibial osteotomy.
J Orthop Sci. 2007;12:214-218.
- 45, Todo M, Nagamine R, and Yamaguchi S.
Stress analysis of PS type knee prosthesis under deep flexion.
Journal of Biomechanical Science and Engineering 2007;2:237-245

- 46, Nagamine R, Kondo K, Nomura H, Kanekasu K, Sonohata M, and Sugioka Y.
Shape of the joint gap for 90 degrees and 120 degrees knee flexion after total
knee arthroplasty. J Orthop Sci. 2008;13:354-358
- 47, Nagamine R, Chen W, Hara T, Kondo K, Sugioka Y
Immediate reduction of white blood cell count after Tocilizumab
administration was observed in some cases
Mod. Rheumatol. 2009;19:348-350
- 48, Nakashima Y, Kondo M, Harada H, Horiuchi T, Ishinishi T, Jojima H, Kuroda K,
Miyahara H, Nagamine R, Nakashima H, Otsuka T, Saikawa I, Shono E, Suematsu E,
Tsuru T, Wada K and Iwamoto Y.
Clinical evaluation of tocilizumab for patients with active rheumatoid arthritis
refractory to anti-TNF biologics: tocilizumab in combination with methotrexate.
Mod Rheumatol. 2010; 20(4):343-52
- 49, Chen W, Nagamine R, Kondo K and Todo M.
Medial Soft Tissue Release is more Effective in Knee Flexion than in
Extension for Medial Osteoarthritic Knees during Posterior Stabilized Total
Knee Arthroplasty
J Orthopaedic Surgery. 2011;19:230-233
- 50, Nakashima Y, Kondo M, Fukuda T, Harada H, Horiuchi T, Ishinishi T, Jojima H,
Kuroda K, Miyahara H, Maekawa M, Nishizaka H, Nagamine R, Nakashima H, Otsuka
T, Shono E, Suematsu E, Shimauchi T, Tsuru T, Wada K, Yoshizawa S, Yoshizawa S,
Iwamoto Y
Remission in patients with active rheumatoid arthritis by tocilizumab treatment
in routine clinical practice: results from 3 years of prospectively registered data.
Mod Rheumatol. 2014;24:258-264
- 51, Osano K, Nagamine R, Todo M, Kawasaki K
The Effect of Malrotation of Tibial Component of Total Knee Arthroplasty on
Tibial Insert during High Flexion Using a Finite Element Analysis
The Scientific World Journal 2014: 695028.
- 52, Mohd Afzan Mohd Anuar, Todo M, Nagamine R and Hirokawa S
Dynamic Finite Element Analysis of Mobile Bearing Type Knee
Prosthesis under Deep Flexional Motion
The Scientific World Journal 2014: 586921

- 53, Takeuchi T, Tanaka Y, Yamanaka H, Amano K, Nagamine R, Park W, Shiozawa K, Tsukano M, Wei JC, Shao J, Togo O, Mashimo H.
Efficacy and safety of olokizumab in Asian patients with moderate-to-severe rheumatoid arthritis, previously exposed to anti-TNF therapy: Results from a randomized phase II trial.
Mod Rheumatol. 2016;26(1):15-23.
- 54, Ma Y, Nagamine R, Chen W.
Comparative evaluation of posterior cruciate ligament in total knee arthroplasty: an in vivo study
J Orthopaedic Surgery (Hongkong). 2017;25:1-6
- 54, Nakashima Y, Kondo M, Shono E, Ishinishi T, Tsukamoto H, Kuroda K, Maeyama A, Harada H, Maekawa M, Shimauchi T, Nagamine R, Jojima H, Yoshizawa S, Tsuru T, Otsuka T, Miyahara H, Suematsu E, Wada K, Yoshizawa S, Inoue Y, Fukuda T, Ikemura S, Haraguchi A.
Suppression of joint destruction with subcutaneous tocilizumab for Japanese patients with rheumatoid arthritis in clinical practice.
Mod Rheumatol. 2019 Oct 17:1-9
- 55, Kawasaki M, Nagamine R, Chen W, Ma Y, Sakai A, Suguro T.
Proximal tibia vara involves the medial shift of the tibial articular surface
J Orthopaedic Surgery (Hongkong). 2020;28(1):2309499020902592.
- 56, Nakashima Y, Kondo M, Shono E, Ishinishi T, Tsukamoto H, Kuroda K, Maeyama A, Harada H, Maekawa M, Shimauchi T, Nagamine R, Jojima H, Yoshizawa S, Tsuru T, Otsuka T, Miyahara H, Suematsu E, Wada K, Yoshizawa S, Inoue Y, Fukuda T, Ikemura S, Haraguchi A.
Suppression of joint destruction with subcutaneous tocilizumab for Japanese patients with rheumatoid arthritis in clinical practice. Mod Rheumatol. 2020 Sep;30(5):807-815. doi: 10.1080/14397595.2019.1676369. Epub 2019 Oct 17.
- 57, Nagamine R, Kawasaki M, Kim KI, Sakai A, Suguro T.
The posterior tibial slope is mainly created by the posterior rotation of the tibial condyles. J Orthop Surg (Hong Kong). 2020 Sep-Dec;28(3): 2309499020975580. doi: 10.1177/2309499020975580
- 58, Kawasaki M, Kaneyama R, Suzuki H, Fujitani T, Tsukamoto M, Sabanai K, Yoshioka T, Okimoto N, Nagamine R, Sakai A.
Component gap control during posterior-stabilised total knee arthroplasty using the posterior condylar pre-cut technique.
J Exp Orthop. 2021 Sep 15;8(1):77